

Preprint, arXiv:0910.4170

SOME q -CONGRUENCES RELATED TO 3-ADIC VALUATIONS

HAO PAN AND ZHI-WEI SUN

Department of Mathematics, Nanjing University
Nanjing 210093, People's Republic of China
haopan79@yahoo.com.cn, zwsun@nju.edu.cn

ABSTRACT. In 1992 Strauss, Shallit and Zagier proved that for any positive integer a we have

$$\sum_{k=0}^{3^a-1} \binom{2k}{k} \equiv 0 \pmod{3^{2a}}$$

and furthermore

$$\frac{1}{3^{2a}} \sum_{k=0}^{3^a-1} \binom{2k}{k} \equiv 1 \pmod{3}.$$

Recently a q -analogue of the former congruence was conjectured by Guo and Zeng. In this paper we prove the conjecture of Guo and Zeng, and also give a q -analogue of the latter congruence.

1. INTRODUCTION

Partially motivated by the work of Pan and Sun [PS], Sun and Tauraso [ST1] proved that for any prime p and $a \in \mathbb{Z}^+ = \{1, 2, 3, \dots\}$ we have

$$\sum_{k=0}^{p^a-1} \binom{2k}{k} \equiv \left(\frac{p^a}{3}\right) \pmod{p^2},$$

where $(-)$ denotes the Legendre symbol. (See also [ST2, ZPS, S09a, S09b, S09c] for related results.) When checking whether there are composite numbers n such that

$$\sum_{k=0}^{n-1} \binom{2k}{k} \equiv \left(\frac{n}{3}\right) \pmod{n^2},$$

2010 *Mathematics Subject Classification*. Primary 11B65; Secondary 05A10, 05A30, 11A07, 11S99.

Keywords: 3-adic valuation, central binomial coefficient, congruence, q -analogue.

The second author is the corresponding author. He was supported by the National Natural Science Foundation (grant 10871087) and the Overseas Cooperation Fund (grant 10928101) of China.

Sun and Tauraso found that

$$\nu_3\left(\sum_{k=0}^{3^a-1} \binom{2k}{k}\right) \geq 2a \quad \text{for } a = 1, 2, 3, \dots, \quad (1.1)$$

where $\nu_3(m)$ denotes the 3-adic valuation of an integer m (i.e., $\nu_3(m) = \sup\{a \in \mathbb{N} : 3^a \mid m\}$ with $\mathbb{N} = \{0, 1, 2, \dots\}$). However, a refinement of this was proved earlier by Strauss, Shallit and Zagier [SSZ] in 1992.

Theorem 1.1 (Strauss, Shallit and Zagier [SSZ]). *For any $a \in \mathbb{Z}^+$ we have*

$$\sum_{k=0}^{3^a-1} \binom{2k}{k} \equiv 3^{2a} \pmod{3^{2a+1}}. \quad (1.2)$$

Furthermore,

$$\frac{\sum_{k=0}^{n-1} \binom{2k}{k}}{n^2 \binom{2n}{n}} \equiv -1 \pmod{3} \quad \text{for all } n \in \mathbb{Z}^+.$$

Recall that the usual q -analogue of $n \in \mathbb{N}$ is

$$[n]_q = \frac{1 - q^n}{1 - q} = \sum_{0 \leq k < n} q^k$$

which tends to n as $q \rightarrow 1$. For $d \in \mathbb{Z}^+$ the d -th cyclotomic polynomial is given by

$$\Phi_d(q) = \prod_{\substack{r=1 \\ (r,d)=1}}^d \left(q - e^{2\pi i r/d} \right) \in \mathbb{Z}[q].$$

Given a positive integer $n > 1$ we obviously have

$$[n]_q = \frac{q^n - 1}{q - 1} = \prod_{k=1}^{n-1} \left(q - e^{2\pi i k/n} \right) = \prod_{\substack{d|n \\ d>1}} \Phi_d(q).$$

It is well known that if $d_1, d_2 \in \mathbb{Z}^+$ are distinct then $\Phi_{d_1}(q)$ and $\Phi_{d_2}(q)$ are relatively prime in the polynomial ring $\mathbb{Z}[q]$. If p is a prime and a is a positive integer, then

$$\Phi_{p^a}(q) = \frac{q^{p^a} - 1}{q^{p^{a-1}} - 1} = [p]_{q^{p^{a-1}}} \quad \text{and} \quad [p^a]_q = \prod_{j=1}^a \Phi_{p^j}(q).$$

For $n, k \in \mathbb{N}$ the usual q -analogue of the binomial coefficient $\binom{n}{k}$ is the following q -binomial coefficient:

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \begin{cases} ([n]_q \cdots [n-k+1]_q) / ([1]_q \cdots [k]_q) & \text{if } 0 < k \leq n, \\ 1 & \text{if } k = 0, \\ 0 & \text{if } k > n. \end{cases}$$

Note that $\begin{bmatrix} n \\ k \end{bmatrix}_q \rightarrow \binom{n}{k}$ as $q \rightarrow 1$. Many combinatorial identities and congruences involving binomial coefficients have their q -analogues (cf. [St]).

Recently Guo and Zeng [GZ] proposed a q -analogue of (1.1), namely they formulated the following conjecture.

Conjecture 1.2 (Guo and Zeng [GZ, Conjecture 3.5]). *Let a be a positive integer. Then*

$$\sum_{k=0}^{3^a m - 1} q^k \begin{bmatrix} 2k \\ k \end{bmatrix}_q \equiv 0 \pmod{[3^a]_q^2} \quad \text{for any } m \in \mathbb{Z}^+. \quad (1.3)$$

Concerning this conjecture, Guo and Zeng [GZ] were able to show (1.3) with the modulus $[3^a]_q^2$ replaced by $[3^a]_q$.

In this paper we confirm Conjecture 1.2 and give a q -analogue of (1.2).

Theorem 1.3. *Let $a \in \mathbb{Z}^+$. Then (1.3) holds. Furthermore, we have the following q -analogue of (1.2):*

$$\frac{1}{[3^a]_q^2} \sum_{k=0}^{3^a - 1} q^k \begin{bmatrix} 2k \\ k \end{bmatrix}_q \equiv 2R(a, q) \pmod{\Phi_{3^a}(q)} \quad (1.4)$$

where

$$R(a, q) := \sum_{\substack{k=1 \\ 3|k-1}}^{3^a-1} q^{\frac{(k+2)(k-1)}{6}} \frac{(-1)^k}{[k]_q^2} \left(1 + \left(\frac{k-1}{3} - \frac{3^{a-1}+1}{2} \right) (1 - q^k) \right). \quad (1.5)$$

Remark 1.4. Let $a \in \mathbb{Z}^+$. Then $\lim_{q \rightarrow 1} R(a, q) \equiv -1 \pmod{3}$ since

$$\sum_{\substack{k=1 \\ 3|k-1}}^{3^a-1} \frac{(-1)^k}{k^2} = \sum_{j=0}^{3^{a-1}-1} \frac{(-1)^{3j+1}}{(3j+1)^2} \equiv - \sum_{j=0}^{3^{a-1}-1} (-1)^j = -1 \pmod{3}.$$

Also, for $k \in \mathbb{Z}^+$ with $k \equiv 1 \pmod{3}$, $[k]_q$ is relatively prime to $[3^a]_q$ since k is relatively prime to 3^a . Therefore (1.4) implies both (1.2) and (1.3) in the case $m = 1$.

We are going to prove an auxiliary result in the next section and then show Theorem 1.3 in Section 3.

2. AN AUXILIARY THEOREM

Theorem 2.1. *Let $a, m \in \mathbb{Z}^+$ and let $\psi : \mathbb{Z} \rightarrow \mathbb{Z}$ be a function such that for any $k \in \mathbb{Z}$ and $j = 1, \dots, a$ we have*

$$\psi(k) \equiv \psi(-k) \pmod{3^a} \quad \text{and} \quad \psi(k + 3^j) \equiv \psi(k) \pmod{3^j}.$$

Then

$$\sum_{k=1}^{3^a m - 1} q^{\psi(k)} \left(\frac{k}{3} \right) \left[\begin{matrix} 2 \cdot 3^a m \\ k \end{matrix} \right]_q \equiv 0 \pmod{[3^a]_q^2}. \quad (2.1)$$

In particular,

$$\sum_{k=1}^{3^a m - 1} \left(\frac{k}{3} \right) \left[\begin{matrix} 2 \cdot 3^a m \\ k \end{matrix} \right]_q \equiv 0 \pmod{[3^a]_q^2}. \quad (2.2)$$

Proof. Clearly $[x]_q \equiv [y]_q \pmod{\Phi_d(q)}$ provided that $x \equiv y \pmod{d}$. By the q -Lucas congruence (cf. [Sa]),

$$\left[\begin{matrix} x_1 d + y_1 \\ x_2 d + y_2 \end{matrix} \right]_q \equiv \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \left[\begin{matrix} y_1 \\ y_2 \end{matrix} \right]_q \pmod{\Phi_d(q)}$$

for $x_1, x_2, y_1, y_2 \in \mathbb{N}$ with $0 \leq y_1, y_2 \leq d - 1$. Recall that

$$[3^a]_q = \prod_{j=1}^a \Phi_{3^j}(q).$$

Since these $\Phi_{3^j}(q)$ are relatively prime and $[2 \cdot 3^a m]_q \equiv 0 \pmod{[3^a]_q}$, we only need to show that

$$\sum_{k=1}^{3^a m - 1} \left(\frac{k}{3} \right) \frac{q^{\psi(k)}}{[k]_q} \left[\begin{matrix} 2 \cdot 3^a m - 1 \\ k - 1 \end{matrix} \right]_q \equiv 0 \pmod{\Phi_{3^j}(q)}$$

for every $j = 1, \dots, a$.

For any $1 \leq j \leq a$ and $1 \leq k \leq 3^a m - 1$ with $3 \nmid k$, write $k = 3^j s + t$ where $1 \leq t \leq 3^j - 1$. Then, by the q -Lucas congruence,

$$\left[\begin{matrix} 2 \cdot 3^a m - 1 \\ k - 1 \end{matrix} \right]_q \equiv \begin{pmatrix} 2 \cdot 3^{a-j} m - 1 \\ s \end{pmatrix} \left[\begin{matrix} 3^j - 1 \\ t - 1 \end{matrix} \right]_q \pmod{\Phi_{3^j}(q)}.$$

And we have

$$\begin{aligned} \left[\begin{matrix} 3^j - 1 \\ t - 1 \end{matrix} \right]_q &= \prod_{j=1}^{t-1} \frac{[3^j - j]_q}{[j]_q} \\ &= \prod_{j=1}^{t-1} \frac{q^{-j}([3^j]_q - [j]_q)}{[j]_q} \equiv (-1)^{t-1} q^{-\binom{t}{2}} \pmod{\Phi_{3^j}(q)}. \end{aligned}$$

Hence

$$\begin{aligned}
& \sum_{k=1}^{3^a m-1} \left(\frac{k}{3} \right) \frac{q^{\psi(k)}}{[k]_q} \left[\begin{matrix} 2 \cdot 3^a m - 1 \\ k - 1 \end{matrix} \right]_q \\
&= \sum_{s=0}^{3^{a-j} m-1} \sum_{t=1}^{3^j-1} \left(\frac{3^j s + t}{3} \right) \frac{q^{\psi(3^j s + t)}}{[3^j s + t]_q} \left[\begin{matrix} 2 \cdot 3^a m - 1 \\ 3^j s + t - 1 \end{matrix} \right]_q \\
&\equiv \sum_{s=0}^{3^{a-j} m-1} \binom{2 \cdot 3^{a-j} m - 1}{s} \sum_{t=1}^{3^j-1} \left(\frac{t}{3} \right) \frac{(-1)^{t-1} q^{\psi(t) - \binom{t}{2}}}{[t]_q} \pmod{\Phi_{3^j}(q)}.
\end{aligned}$$

Clearly,

$$\begin{aligned}
& 2 \sum_{t=1}^{3^j-1} \left(\frac{t}{3} \right) \frac{(-1)^{t-1} q^{\psi(t) - \binom{t}{2}}}{[t]_q} \\
&= \sum_{t=1}^{3^j-1} \left(\left(\frac{t}{3} \right) \frac{(-1)^{t-1} q^{\psi(t) - \binom{t}{2}}}{[t]_q} + \left(\frac{3^j - t}{3} \right) \frac{(-1)^{3^j-t-1} q^{\psi(3^j-t) - \binom{3^j-t}{2}}}{[3^j - t]_q} \right) \\
&\equiv \sum_{\substack{t=1 \\ 3 \nmid t}}^{3^j-1} \left(\frac{t}{3} \right) \left(\frac{(-1)^{t-1} q^{\psi(t) - \binom{t}{2}}}{[t]_q} + \frac{(-1)^{t-1} q^{\psi(t) - \binom{-t}{2}}}{-q^{-t} [t]_q} \right) = 0 \pmod{\Phi_{3^j}(q)}.
\end{aligned}$$

So (2.1) holds.

Note that (2.2) is just (2.1) with ψ replaced by the zero function from $\mathbb{Z} \rightarrow \mathbb{Z}$. So (2.2) is also valid. This concludes the proof. $\square$

3. PROOF OF THEOREM 1.3

Lemma 3.1. *Suppose that $k \equiv l \pmod{3^a}$ where $k, l \in \mathbb{Z}$ and $a \in \mathbb{Z}^+$. Then*

$$2k^2 - k \left(\frac{k}{3} \right) \equiv 2l^2 - l \left(\frac{l}{3} \right) \pmod{3^{a+1}}.$$

Proof. Observe that

$$2(k+l) - \left(\frac{k}{3} \right) \equiv 4k - \left(\frac{k}{3} \right) \equiv 0 \pmod{3}.$$

Thus

$$\begin{aligned}
& 2k^2 - k \left(\frac{k}{3} \right) - \left(2l^2 - l \left(\frac{l}{3} \right) \right) \\
&= (k-l) \left(2(k+l) - \left(\frac{k}{3} \right) \right) \equiv 0 \pmod{3^{a+1}}.
\end{aligned}$$

We are done. $\square$

Lemma 3.2. *Let $a \in \mathbb{Z}^+$ and let ψ be a function as in Theorem 2.1. Then*

$$\begin{aligned} & \frac{1}{2[3^a]_q^2} \sum_{k=1}^{3^a-1} q^{\psi(k)} \left(\frac{k}{3}\right) \begin{bmatrix} 2 \cdot 3^a \\ k \end{bmatrix}_q \\ & \equiv \sum_{\substack{k=1 \\ 3 \nmid k-1}}^{3^a-1} q^{\psi(k)-\binom{k}{2}} \frac{(-1)^{k-1}}{[k]_q^2} (1 + \Psi_a(k)(1 - q^k)) \pmod{\Phi_{3^a}(q)}, \end{aligned} \quad (3.1)$$

where

$$\Psi_a(k) := \frac{\psi(3^a - k) - \psi(k)}{3^a} + \frac{3^a - 1}{2} - k. \quad (3.2)$$

Proof. We have

$$\begin{aligned} & \sum_{k=1}^{3^a-1} \left(\frac{k}{3}\right) \frac{q^{\psi(k)}}{[k]_q} \begin{bmatrix} 2 \cdot 3^a - 1 \\ k - 1 \end{bmatrix}_q \\ & = \sum_{k=1}^{3^a-1} \left(\frac{k}{3}\right) \frac{q^{\psi(k)}}{[k]_q} \prod_{j=1}^{k-1} \frac{q^{-j}([2 \cdot 3^a]_q - [j]_q)}{[j]_q} \\ & \equiv \sum_{k=1}^{3^a-1} \left(\frac{k}{3}\right) \frac{(-1)^{k-1} q^{\psi(k)-\binom{k}{2}}}{[k]_q} \left(1 - 2 \sum_{j=1}^{k-1} \frac{[3^a]_q}{[j]_q}\right) \pmod{\Phi_{3^a}(q)^2}, \end{aligned}$$

since

$$[2 \cdot 3^a]_q = [3^a]_q(1 + q^{3^a}) = [3^a]_q(2 + q^{3^a} - 1) \equiv 2[3^a]_q \pmod{[3^a]_q^2}.$$

Note that for $s = 0, 1, 2, \dots$ we have

$$\begin{aligned} q^{3^a s} &= 1 + (q^{3^a} - 1) \sum_{j=0}^{s-1} q^{3^a j} = 1 + (q^{3^a} - 1) \left(s + \sum_{j=0}^{s-1} (q^{3^a j} - 1)\right) \\ &\equiv 1 + s(q^{3^a} - 1) \pmod{\Phi_{3^a}(q)^2} \end{aligned}$$

and

$$q^{-3^a s} \equiv \frac{1}{1 + s(q^{3^a} - 1)} = \frac{1 - s(q^{3^a} - 1)}{1 - s^2(q^{3^a} - 1)^2} \equiv 1 - s(q^{3^a} - 1) \pmod{\Phi_{3^a}(q)^2}.$$

Also, for each $1 \leq k \leq 3^a - 1$, we have

$$\begin{aligned}
& \frac{q^{\psi(3^a-k)-\binom{3^a-k}{2}}}{[3^a-k]_q} \left(1 - 2 \sum_{j=1}^{3^a-k-1} \frac{[3^a]_q}{[j]_q} \right) \\
&= \frac{q^{\psi(3^a-k)-\binom{3^a}{2}+3^a k-\binom{k+1}{2}} ([3^a]_q + [k]_q)}{q^{-k}([3^a]_q^2 - [k]_q^2)} \left(1 - 2 \sum_{j=k+1}^{3^a-1} \frac{[3^a]_q}{[3^a-j]_q} \right) \\
&\equiv \frac{q^{\psi(k)-\binom{k}{2}} (1 + (\frac{\psi(3^a-k)-\psi(k)}{3^a} + k - \frac{3^a-1}{2})(q^{3^a}-1))([3^a]_q + [k]_q)}{-[k]_q^2} \\
&\quad \times \left(1 + 2 \sum_{j=k+1}^{3^a-1} \frac{q^j [3^a]_q}{[j]_q} \right) \\
&\equiv - \frac{q^{\psi(k)-\binom{k}{2}}}{[k]_q} \left(1 + \left(\frac{\psi(3^a-k)-\psi(k)}{3^a} + k - \frac{3^a-1}{2} \right) (q^{3^a}-1) \right) \\
&\quad - 2 \frac{q^{\psi(k)-\binom{k}{2}}}{[k]_q} \sum_{j=k+1}^{3^a-1} \frac{q^j [3^a]_q}{[j]_q} - \frac{q^{\psi(k)-\binom{k}{2}} [3^a]_q}{[k]_q^2} \pmod{\Phi_{3^a}(q)^2}.
\end{aligned}$$

Clearly,

$$\sum_{j=k+1}^{3^a-1} \frac{q^j}{[j]_q} = \sum_{j=k+1}^{3^a-1} \frac{1+q^j-1}{[j]_q} = -(3^a-1-k)(1-q) + \sum_{j=k+1}^{3^a-1} \frac{1}{[j]_q},$$

and

$$\begin{aligned}
& \sum_{j=1}^{3^a-1} \frac{1}{[j]_q} = \frac{1}{2} \sum_{j=1}^{3^a-1} \left(\frac{1}{[j]_q} + \frac{1}{[3^a-j]_q} \right) \\
&\equiv \frac{1}{2} \sum_{j=1}^{3^a-1} \left(\frac{1}{[j]_q} - \frac{q^j}{[j]_q} \right) = \frac{3^a-1}{2} (1-q) \pmod{\Phi_{3^a}(q)}.
\end{aligned}$$

Thus we get

$$\begin{aligned}
& \frac{q^{\psi(k)-\binom{k}{2}}}{[k]_q} \left(1 - \sum_{j=1}^{k-1} \frac{[2 \cdot 3^a]_q}{[j]_q} \right) + \frac{q^{\psi(3^a-k)-\binom{3^a-k}{2}}}{[3^a-k]_q} \left(1 - \sum_{j=1}^{3^a-k-1} \frac{[2 \cdot 3^a]_q}{[j]_q} \right) \\
&\equiv - \frac{q^{\psi(k)-\binom{k}{2}}}{[k]_q} \left(\left(\frac{\psi(3^a-k)-\psi(k)}{3^a} + \frac{3}{2}(3^a-1)-k \right) (q^{3^a}-1) \right) \\
&\quad - \frac{q^{\psi(k)-\binom{k}{2}}}{[k]_q} \left(2 \sum_{j=1}^{k-1} \frac{[3^a]_q}{[j]_q} + 2 \sum_{j=k+1}^{3^a-1} \frac{[3^a]_q}{[j]_q} + \frac{q^{\psi(k)-\binom{k}{2}} [3^a]_q}{[k]_q^2} \right) \\
&\equiv \frac{q^{\psi(k)-\binom{k}{2}} [3^a]_q}{[k]_q^2} + \frac{q^{\psi(k)-\binom{k}{2}}}{[k]_q} \Psi_a(k) (1-q^{3^a}) \pmod{\Phi_{3^a}(q)^2}.
\end{aligned}$$

It follows that

$$\begin{aligned} & \sum_{k=1}^{3^a-1} \left(\frac{k}{3}\right) \frac{q^{\psi(k)}}{[k]_q} \begin{bmatrix} 2 \cdot 3^a - 1 \\ k - 1 \end{bmatrix}_q \\ & \equiv \sum_{\substack{k=1 \\ 3 \nmid k-1}}^{3^a-1} (-1)^{k-1} \frac{q^{\psi(k)-\binom{k}{2}}}{[k]_q} \left(\frac{[3^a]_q}{[k]_q} + \Psi_a(k)(1 - q^{3^a}) \right) \pmod{\Phi_{3^a}(q)^2}. \end{aligned}$$

Noting that $[3^a]_q$ divides both sides of the above congruence by Theorem 2.1 and $[2 \cdot 3^a] \equiv 2[3^a] \pmod{[3^a]_q^2}$, we are done. $\square$

Proof of Theorem 1.3. Let $m \in \mathbb{Z}^+$. By [T, (4.3)] in the case $d = 0$, we have

$$\sum_{k=0}^{3^a m - 1} q^k \begin{bmatrix} 2k \\ k \end{bmatrix}_q = - \sum_{k=1}^{3^a m - 1} q^{\psi_m(k)} \left(\frac{k}{3}\right) \begin{bmatrix} 2 \cdot 3^a m \\ k \end{bmatrix}_q, \quad (3.3)$$

where

$$\psi_m(k) = \frac{2(3^a m - k)^2 - (3^a m - k) \left(\frac{3^a m - k}{3}\right) - 1}{3}.$$

According to Lemma 3.1, the function $\psi = \psi_m$ has the property described in Theorem 2.1. Combining (2.1) with (3.3) we get (1.3).

Now it remains to prove (1.4). By (3.3) and Lemma 3.2, we finally obtain

$$\begin{aligned} & \sum_{k=0}^{3^a-1} q^k \begin{bmatrix} 2k \\ k \end{bmatrix}_q = - \sum_{k=1}^{3^a-1} q^{\psi_1(k)} \left(\frac{k}{3}\right) \begin{bmatrix} 2 \cdot 3^a \\ k \end{bmatrix}_q \\ & \equiv 2[3^a]_q^2 \sum_{\substack{k=1 \\ 3 \nmid k-1}}^{3^a-1} q^{\frac{(k+2)(k-1)}{6}} \frac{(-1)^k}{[k]_q^2} \left(1 + \left(\frac{k-1}{3} - \frac{3^{a-1}+1}{2} \right) (1 - q^k) \right) \\ & \pmod{\Phi_{3^a}(q)[3^a]_q^2}. \end{aligned}$$

This concludes our proof. $\square$

Acknowledgment. The authors are indebted to Prof. J. Shallit for informing the reference [SSZ].

REFERENCES

- [GZ] V. J. W. Guo and J. Zeng, *Some congruences involving central q -binomial coefficients*, preprint, arXiv:0910.3563. <http://arxiv.org/abs/0910.3563>.
- [PS] H. Pan and Z. W. Sun, *A combinatorial identity with application to Catalan numbers*, Discrete Math. **306** (2006), 1921–1940.
- [Sa] B. E. Sagan, *Congruence properties of q -analogs*, Adv. in Math. **95** (1992), 127–143.

- [SSZ] N. Strauss, J. Shallit and D. Zagier, *Some strange 3-adic identities*, Amer. Math. Monthly **99** (1992), 66–69.
- [St] R. P. Stanley, *Enumerative Combinatorics*, Vol. 2, Cambridge Univ. Press, Cambridge, 1999.
- [S06] Z. W. Sun, *Binomial coefficients and quadratic fields*, Proc. Amer. Math. Soc. **134** (2006), 2213–2222.
- [S09a] Z. W. Sun, *Binomial coefficients, Catalan numbers and Lucas quotients*, preprint, arXiv:0909.5648. <http://arxiv.org/abs/0909.5648>.
- [S09b] Z. W. Sun, *Various congruences involving binomial coefficients and higher-order Catalan numbers*, preprint, arXiv:0909.3808. <http://arxiv.org/abs/0909.3808>.
- [S09c] Z. W. Sun, *p -adic valuations of some sums of multinomial coefficients*, preprint, arXiv:0910.3892. <http://arxiv.org/abs/0910.3892>.
- [ST1] Z. W. Sun and R. Tauraso, *On some new congruences for binomial coefficients*, Acta Arith., to appear. <http://arxiv.org/abs/0709.1665>.
- [ST2] Z. W. Sun and R. Tauraso, *New congruences for central binomial coefficients*, Adv. in Math., to appear. <http://arxiv.org/abs/0805.0563>.
- [T] R. Tauraso, *q -analogs of some congruences involving Catalan numbers*, preprint, arXiv:0905.3816. <http://arxiv.org/abs/0905.3816>.
- [ZPS] L. Zhao, H. Pan and Z. W. Sun, *Some congruences for the second-order Catalan numbers*, Proc. Amer. Math. Soc. **138** (2010), 37–46.